

GLOBAL GORENSTEIN DIMENSIONS OF POLYNOMIAL RINGS AND OF DIRECT PRODUCT OF RINGS

D. BENNIS AND N. MAHDOU

ABSTRACT. In this paper, we extend the well-known Hilbert's syzygy theorem to the Gorenstein homological dimensions of rings. Also, we study the Gorenstein homological dimensions of direct product of rings, which gives examples of non-Noetherian rings of finite Gorenstein dimensions and infinite classical weak dimension.

1. INTRODUCTION

Throughout this paper all rings are commutative with identity element and all modules are unital.

Setup and Notation: Let R be a ring, and let M be an R -module.

As usual we use $\text{pd}_R(M)$, $\text{id}_R(M)$, and $\text{fd}_R(M)$ to denote, respectively, the classical projective, injective and flat dimensions of M . $\text{gldim}(R)$ and $\text{wdim}(R)$ are, respectively, the classical global and weak dimensions of R .

It is by now a well-established fact that even if R to be non-Noetherian, there exist Gorenstein projective, injective and flat dimensions of M , which are usually denoted by $\text{Gpd}_R(M)$, $\text{Gid}_R(M)$, and $\text{Gfd}_R(M)$, respectively. Some references are [1, 2, 4, 5, 7].

Recently, the authors in [2] started the study of global Gorenstein dimensions of rings, which are called, for a ring R , Gorenstein projective, injective, and weak dimensions of R , denoted by $\text{GPD}(R)$, $\text{GID}(R)$, and $G - \text{wdim}(R)$, respectively, and, respectively, defined as follows:

$$\begin{aligned}\text{GPD}(R) &= \sup\{\text{Gpd}_R(M) \mid M \text{ } R\text{-module}\}, \\ \text{GID}(R) &= \sup\{\text{Gid}_R(M) \mid M \text{ } R\text{-module}\}, \text{ and} \\ G - \text{wdim}(R) &= \sup\{\text{Gfd}_R(M) \mid M \text{ } R\text{-module}\}.\end{aligned}$$

They proved that, for any ring R : $G - \text{wdim}(R) \leq \text{GID}(R) = \text{GPD}(R)$ [2, Theorems 3.2 and 4.2]. So, according to the terminology of the classical theory of homological dimensions of rings, the common value of $\text{GPD}(R)$ and $\text{GID}(R)$ is called *Gorenstein global dimension* of R , and denoted by $\mathbf{G-gldim}(R)$.

They also proved that the Gorenstein global and weak dimensions are refinements of the classical global and weak dimensions of rings, respectively. That is [2,

Key words and phrases. Gorenstein homological dimensions of modules and of rings; strongly Gorenstein projective, injective, and flat modules; $(n-)$ Gorenstein rings; quasi-Frobenius rings; n -FC rings; IF-rings (weakly quasi-Frobenius rings); Hilbert's syzygy theorem; direct product of rings.

Propositions 3.11 and 4.5]: $G\text{-gldim}(R) \leq \text{gldim}(R)$ and $G\text{-wdim}(R) \leq \text{wdim}(R)$, with equality if $\text{wdim}(R)$ is finite.

In Section 2, we are mainly interested in computing the Gorenstein dimensions of commutative polynomial rings to generalize the fact that a polynomial ring over Gorenstein ring is also Gorenstein. Explicitly, we extent the equalities of the well-known Hilbert's syzygy theorem to the Gorenstein global and weak dimensions (please see Theorems 2.1 and 2.11).

Recall that a ring R is said to be n -Gorenstein, for a positive integer n , if it is Noetherian and $\text{id}_R(R) \leq n$ [8]. And R is said to be Gorenstein, if it is n -Gorenstein for some positive integer n . These kind of rings have a characterization by the weak and global Gorenstein dimensions of rings. Namely, we have [2, Corollary 2.3]: If R is a Noetherian ring, then $G\text{-wdim}(R) = G\text{-gldim}(R)$, such that, for a positive integer n , $G\text{-gldim}(R) \leq n$ if, and only if, R is n -Gorenstein. In the case $n = 0$, we have the last equivalence without assuming in the first place that R to be Noetherian; that is [2, Theorem 5.2.4]: $G\text{-gldim}(R) = 0$ if, and only if, R is 0-Gorenstein (i.e., quasi-Frobenius). Thus, the study of the Gorenstein dimensions of polynomial rings allows us to give examples of Noetherian rings which have finite Gorenstein global (=weak) dimension and infinite global (=weak) dimension (see Example 2.10). Obviously, one would like to have examples out of the class of Noetherian rings. Explicitly, one would like to have examples of a family of non-Noetherian rings $\{R_i\}_{i \geq 1}$ such that $G\text{-wdim}(R_i) = i$, $G\text{-gldim}(R_i) > i$, and $\text{wdim}(R_i) = \infty$ for all $i \geq 1$. This will be obtained in Section 3 (Example 3.8) after a study of the Gorenstein dimensions of direct product of rings (Theorems 3.1 and 3.5).

2. GORENSTEIN DIMENSIONS OF POLYNOMIAL RINGS

The aim of this section is to compute the Gorenstein global and weak dimensions of commutative polynomial rings.

Through this section, R denotes a commutative ring, and $R[X]$ denotes the polynomial ring in one indeterminate over R . We use $M[X]$, where M is an R -module, to denote the $R[X]$ -module $M \otimes_R R[X]$.

The first main result in this section is:

Theorem 2.1. *Let $R[X_1, X_2, \dots, X_n]$ be the polynomial ring in n indeterminates over R . Then: $G\text{-gldim}(R[X_1, X_2, \dots, X_n]) = G\text{-gldim}(R) + n$.*

To prove this theorem, we need some results.

First, we need the notion of strongly Gorenstein projective modules, which are introduced in [1] to characterize the Gorenstein projective modules.

Definition 2.2 ([1], Definition 2.1). *A module M is said to be strongly Gorenstein projective, if there exists an exact sequence of the form: $\mathbf{P} = \dots \xrightarrow{f} P \xrightarrow{f} P \xrightarrow{f} P \xrightarrow{f} \dots$ where P is a projective R -module and f is an endomorphism of P , such that $M \cong \text{Im}(f)$ and such that $\text{Hom}(-, Q)$ leaves the sequence $\mathbf{P}$ exact whenever Q is a projective module.*

These strongly Gorenstein projective modules are a simple characterization, and they are used to characterize the Gorenstein projective modules. We have:

Proposition 2.3 ([2], Proposition 3.9). *A module M is strongly Gorenstein projective if, and only if, there exists a short exact sequence of modules $0 \rightarrow M \rightarrow P \rightarrow M \rightarrow 0$ where P is a projective module, and $\text{Ext}^i(M, Q) = 0$ for some integer $i > 0$ and for any module Q with finite projective dimension (or for any projective module Q).*

Theorem 2.4 ([1], Theorem 2.7). *A module is Gorenstein projective if, and only if, it is a direct summand of a strongly Gorenstein projective module.*

Lemma 2.5. *For any R -module M , $\text{Gpd}_{R[X]}(M[X]) \leq \text{Gpd}_R(M)$.*

Proof. The standard argument shows that the inequality is a simple consequence of the implication: M is a Gorenstein projective R -module $\Rightarrow M[X]$ is a Gorenstein projective $R[X]$ -module. So, assume M to be a Gorenstein projective R -module. We claim that $M[X]$ is a Gorenstein projective $R[X]$ -module.

From Theorem 2.4, M is a direct summand of a strongly Gorenstein projective R -module N . Then, by Proposition 2.3, there exists a short exact sequence of R -modules $0 \rightarrow N \rightarrow P \rightarrow N \rightarrow 0$ where P is a projective R -module and $\text{Ext}_R(N, Q) = 0$ for any projective R -module Q . Hence, we have the following short exact sequence of $R[X]$ -modules $0 \rightarrow N[X] \rightarrow P[X] \rightarrow N[X] \rightarrow 0$ such that $P[X]$ is a projective $R[X]$ -module; and from [11, Theorem 11.65], $\text{Ext}_{R[X]}(N[X], Q) \cong \text{Ext}_R(N, Q) = 0$ for every projective $R[X]$ -module Q (since Q is also projective R -module). Thus, $N[X]$ is a strongly Gorenstein projective $R[X]$ -module (by Proposition 2.3). Therefore, $M[X]$ is a Gorenstein projective $R[X]$ -module as a direct summand of the strongly Gorenstein projective $R[X]$ -module $N[X]$ (by [7, Theorem 2.5]). ■

Lemma 2.6. *For any short exact sequence of modules $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$, we have inequality: $\text{Gpd}(C) \leq 1 + \max\{\text{Gpd}(A), \text{Gpd}(B)\}$.*

Proof. The argument is analogous to the proof of [4, Corollary 1.2.9 (a)]. ■

Lemma 2.7. *Let $R[X]$ be the polynomial ring in one indeterminate over R . Then, $G - \text{gldim}(R[X]) \leq G - \text{gldim}(R) + 1$.*

Proof. Using Lemmas 2.6 and 2.5, the argument is analogous to the proof from [11, Lemma 9.29]. ■

Lemma 2.8. *If all projective $R[X]$ -modules have finite injective dimension, then, for any R -module M , $\text{Gpd}_{R[X]}(M[X]) = \text{Gpd}_R(M)$. In particular, $G - \text{gldim}(R) \leq G - \text{gldim}(R[X])$.*

Proof. From Lemma 2.5, it remains to prove: $\text{Gpd}_{R[X]}(M[X]) \geq \text{Gpd}_R(M)$. The standard argument shows that this inequality follows from the implication: $M[X]$ is a Gorenstein projective $R[X]$ -module $\Rightarrow M$ is a Gorenstein projective R -module. So, assume that $M[X]$ is a Gorenstein projective $R[X]$ -module. We claim that M is a Gorenstein projective R -module.

From Theorem 2.4, $M[X]$ is a direct summand of a strongly Gorenstein projective $R[X]$ -module N . For such $R[X]$ -module N , and from Proposition 2.3, there exists a short exact sequence of $R[X]$ -modules $(\star) 0 \rightarrow N \rightarrow P \rightarrow N \rightarrow 0$ where P is a projective $R[X]$ -module (then it is also projective as an R -module), and $\text{Ext}_{R[X]}(N, Q') = 0$ for every $R[X]$ -module Q' with finite projective dimension. Note that M is a direct summand of N , since it is a direct summand of $M[X]$. Then, to conclude the proof, it suffices, by [7, Theorem 2.5], to prove that N is also a strongly Gorenstein projective R -module. For that, and from the short exact sequence $(\star)$, it remains to prove, from Proposition 2.3, that $\text{Ext}_R^n(N, L) = 0$ for some integer $n > 0$ and for every projective R -module L . From [11, Theorem 11.65], $\text{Ext}_{R[X]}^i(N[X], L) \cong \text{Ext}_R^i(N, L)$ for every $R[X]$ -module L . If L is a projective R -module, then $\text{pd}_{R[X]}(L) = 1$ (by [9, Theorem C, page 124]). Thus, the hypothesis implies that $\text{id}_{R[X]}(L)$ is finite. Then, there exists an integer $n > 0$ such that $\text{Ext}_{R[X]}^n(N[X], L) = 0$. Therefore, $\text{Ext}_R^n(N, L) = 0$, as desired. Finally, $G\text{-gldim}(R) \leq G\text{-gldim}(R[X])$ is clear by above and this completes the proof of Lemma 2.8. ■

Proof of Theorem 2.1. By induction on n , we may assume $n = 1$, such that we write $R[X] = R[X_1]$. Lemmas 2.7 and 2.8 give:

$$G\text{-gldim}(R) \leq G\text{-gldim}(R[X]) \leq G\text{-gldim}(R) + 1.$$

Then, we may assume that $G\text{-gldim}(R)$ and $G\text{-gldim}(R[X])$ are finite. Let $G\text{-gldim}(R) = m$, for some positive integer m . From [2, Lemma 3.3], there exist an R -module M and a projective R -module P such that: $\text{Ext}_R^m(M, P) \neq 0$. Then, from Rees's theorem [11, Theorem 9.37]: $\text{Ext}_{R[X]}^{m+1}(M, P[X]) \cong \text{Ext}_R^m(M, P) \neq 0$, which implies, from [7, Theorem 2.20] and since $P[X]$ is a projective $R[X]$ -module, that $\text{Gpd}_{R[X]}(M) \geq m + 1$. Therefore, $G\text{-gldim}(R[X]) \geq m + 1$, and then the desired equality. ■

Corollary 2.9. *Let $R[X_1, X_2, \dots, X_n, \dots]$ be the polynomial in infinity of indeterminates. Then, $G\text{-gldim}(R[X_1, X_2, \dots, X_n, \dots]) = \infty$.*

Proof. Obvious. ■

Theorem 2.1 allows us to give a family $\{R_i\}_{i \in \mathbb{N}}$ of commutative rings, such that $G\text{-gldim}(R_i) = i$ and $\text{wdim}(R_i) = \infty$ for all $i \in \mathbb{N}$, as shown by the following example:

Example 2.10. *Consider a non-semisimple quasi-Frobenius ring (for example, $K[X]/(X^2)$, where K is a field). Then, for every positive integer n , we have:*

$$G\text{-gldim}(R[X_1, X_2, \dots, X_n]) = n \quad \text{and} \quad \text{wdim}(R[X_1, X_2, \dots, X_n]) = \infty.$$

Proof. We have $G\text{-gldim}(R) = 0$ since R is quasi-Frobenius. On the other hand, $\text{gldim}(R) (= \text{wdim}(R)) = \infty$ by [2, Proposition 3.11] since $\text{gldim}(R) \neq 0$ (since R is a non-semisimple ring) and R is Noetherian (since R is quasi-Frobenius). ■

Now, we treat the Gorenstein weak dimension of polynomial rings.

Theorem 2.11. *If the polynomial ring $R[X]$ in one indeterminate X over R is coherent, then: $G\text{-wdim}(R[X]) = G\text{-wdim}(R) + 1$.*

Proof. First, note that $R = R[X]/(X)$ is also coherent (by [6, Theorem 4.1.1 (1)]), since R is a finitely presented $R[X]$ -module (by the short exact sequence of $R[X]$ -modules $0 \rightarrow R[X] \rightarrow R[X] \rightarrow R \rightarrow 0$), and since $R[X]$ is coherent.

We proceed similarly to the proof of Theorem 2.1. Then, we first show that $G\text{-wdim}(R[X]) \leq G\text{-wdim}(R) + 1$. This inequality is obtained in the same way as the one of Corollary ?? using [7, Proposition 3.10] instead of Lemma 2.5 and the fact that the inequality of Lemma 2.7 remains true, over coherent rings, for the Gorenstein flat dimension case.

Secondly, we must show that $G\text{-wdim}(R) \leq G\text{-wdim}(R[X])$. For that, it suffices to prove the inequality $\text{Gfd}_{R[X]}(M[X]) \geq \text{Gfd}_R(M)$, for every R -module M . Also, the proof of this inequality is similar to the one in the proof of Lemma 2.8, using the properties of strongly Gorenstein flat modules [1, Definition 3.1] ([2, Proposition 4.6] and [1, Theorem 3.5]) instead of the one of the strongly Gorenstein projective modules used in the proof of Lemma 2.8, and using the fact that, over coherent rings, the set of all Gorenstein flat modules is closed under direct summands [7, Theorem 3.7]. And finally, we use [11, Theorem 11.64], [9, Theorem 202]), and [2, Theorem 4.11 (8)] instead of [11, Theorem 11.65], [9, Theorem C, page 124], and [2, Proposition 3.13], respectively.

Now, we have the inequality $G\text{-wdim}(R) \leq G\text{-wdim}(R[X]) \leq G\text{-wdim}(R) + 1$, then we may assume that $G\text{-wdim}(R)$ and $G\text{-wdim}(R[X])$ are finite.

Let $G\text{-wdim}(R) = m$, for some positive integer m . Hence, from [2, Theorem 4.11], there exists a finitely presented R -module M such that $\text{Ext}_R^m(M, R) \neq 0$. Then, from Rees's theorem [11, Theorem 9.37]: $\text{Ext}_{R[X]}^{m+1}(M, R[X]) \cong \text{Ext}_R^m(M, R) \neq 0$. The R -module M is also finitely presented as an $R[X]$ -module (by [6, Theorem 2.1.7], and since R is a finitely presented $R[X]$ -module, by the short exact sequence of $R[X]$ -modules $0 \rightarrow R[X] \rightarrow R[X] \rightarrow R \rightarrow 0$). Then, from [2, Theorem 4.11], $G\text{-wdim}(R[X]) \geq m + 1$, which implies the desired equality. ■

3. GORENSTEIN DIMENSIONS OF DIRECT PRODUCT OF RINGS

The aim of this section is to compute the Gorenstein global and weak dimensions of direct product of commutative rings.

We begin by the Gorenstein global dimension case, which is a generalization of the classical equality: $\text{gldim}(\prod_{i=1}^m R_i) = \sup\{\text{gldim}(R_i), 1 \leq i \leq m\}$ where $\{R_i\}_{i=1, \dots, m}$ is a family of rings [3, Chapter VI, Exercice 8, page 123].

Theorem 3.1. *Let $\{R_i\}_{i=1, \dots, m}$ be a family of rings. Then:*

$$G\text{-gldim}\left(\prod_{i=1}^m R_i\right) = \sup\{G\text{-gldim}(R_i), 1 \leq i \leq m\}.$$

To prove this theorem, we need the following results:

Lemma 3.2. *Let $R \rightarrow S$ be a ring homomorphism such that S is a projective R -module. If M is a (strongly) Gorenstein projective R -module, then $M \otimes_R S$ is a (strongly) Gorenstein projective S -module.*

Namely, we have: $\text{Gpd}_S(M \otimes_R S) \leq \text{Gpd}_R(M)$.

Proof. Assume at first M to be a strongly Gorenstein projective R -module. Then, there exists a short exact sequence of R -modules $0 \rightarrow M \rightarrow P \rightarrow M \rightarrow 0$ where P

is a projective R -module, and $\text{Ext}_R(M, Q) = 0$ for any projective R -module Q . Since S is a projective (then flat) R -module, we have a short exact sequence of S -modules $0 \rightarrow M \otimes_R S \rightarrow P \otimes_R S \rightarrow M \otimes_R S \rightarrow 0$ such that $P \otimes_R S$ is a projective S -module, and for any projective S -module (then projective R -module) L , $\text{Ext}_S(M \otimes_R S, L) = \text{Ext}_R(M, L) = 0$ (by [11, Theorem 11.65]). This implies that $M \otimes_R S$ is a strongly Gorenstein projective S -module.

Now, let M be any arbitrary Gorenstein projective R -module. Then, it is a direct summand of a strongly Gorenstein projective R -module N . Then, $M \otimes_R S$ is a direct summand of the S -module $N \otimes_R S$ which is, from the reason above, strongly Gorenstein projective. Therefore, $M \otimes_R S$ is a Gorenstein projective S -module, as desired. ■

Lemma 3.3. *Let $\{R_i\}_{i=1,\dots,m}$ be a family of rings such that all projective R_i -modules have finite injective dimension, for $i=1,\dots,m$. Let M_i be an R_i -module for $i=1,\dots,m$. If each M_i is a (strongly) Gorenstein projective R_i -module, then $\prod_{i=1}^m M_i$ is a (strongly) Gorenstein projective $(\prod_{i=1}^m R_i)$ -module. Namely, we have: $\text{Gpd}_{(\prod_{i=1}^m R_i)}(\prod_{i=1}^m M_i) \leq \sup\{\text{Gpd}_{R_i}(M_i), 1 \leq i \leq m\}$.*

Proof. By induction on m , it suffices to prove the assertion for $m = 2$.

We assume at first that M_i is a strongly Gorenstein projective R_i -module for $i = 1, 2$. Then, there exists a short exact sequence of R_i -modules $0 \rightarrow M_i \rightarrow P_i \rightarrow M_i \rightarrow 0$ where P_i is a projective R_i -module. Hence, we have a short exact sequence of $R_1 \times R_2$ -modules $0 \rightarrow M_1 \times M_2 \rightarrow P_1 \times P_2 \rightarrow M_1 \times M_2 \rightarrow 0$ where $P_1 \times P_2$ is a projective $R_1 \times R_2$ -module (by [10, Lemma 2.5 (2)]).

On the other hand, let Q be a projective $R_1 \times R_2$ -module. We have:

$$Q = Q \otimes_{R_1 \times R_2} (R_1 \times R_2) = Q \otimes_{R_1 \times R_2} (R_1 \times 0 \oplus 0 \times R_2) = Q_1 \times Q_2$$

where $Q_i = Q \otimes_{R_1 \times R_2} R_i$ for $i = 1, 2$. From [10, Lemma 2.5 (2)], Q_i is a projective R_i -module for $i = 1, 2$. Hence, by hypothesis $\text{id}_{R_i}(Q_i) < \infty$ for $i = 1, 2$, and from [3, Chapter VI, Exercice 10, page 123], $\text{id}_{R_1 \times R_2}(Q_i) \leq \text{id}_{R_i}(Q_i) < \infty$ for $i = 1, 2$. Thus, $\text{id}_{R_1 \times R_2}(Q_1 \times Q_2) < \infty$, so $\text{Ext}_{R_1 \times R_2}^k(M_1 \times M_2, Q_1 \times Q_2) = 0$ for some positive integer k . This implies, from Proposition 2.3, that $M_1 \times M_2$ is a strongly Gorenstein projective $R_1 \times R_2$ -module.

Now, let M_i be any arbitrary Gorenstein projective R_i -module for $i = 1, 2$. Then, there exist an R_i -module G_i and a strongly Gorenstein projective R_i -module N_i for $i = 1, 2$ such that $M_i \oplus G_i = N_i$. Then, $(M_1 \times M_2) \oplus (G_1 \times G_2) = (M_1 \oplus G_1) \times (M_2 \oplus G_2) = N_1 \times N_2$. Since, by the reason above, $N_1 \times N_2$ is a strongly Gorenstein projective $R_1 \times R_2$ -module, and from Theorem 2.4, $M_1 \times M_2$ is a Gorenstein projective $R_1 \times R_2$ -module, as desired. ■

Proof of Theorem 3.1. By induction on m , it suffices to prove the equality for $m = 2$. We must show that, for any positive integer d , we have the equivalence:

$$\text{G-gldim}(R_1 \times R_2) \leq d \iff \text{G-gldim}(R_1) \leq d \text{ and } \text{G-gldim}(R_2) \leq d.$$

Then, assume that $\text{G-gldim}(R_1 \times R_2) \leq d$ for some positive integer d .

Let M_i be an R_i -module for $i = 1, 2$. Since each R_i is a projective $R_1 \times R_2$ -module, and from Lemma 3.2, we have: $\text{Gpd}_{R_i}(M_i) = \text{Gpd}_{R_i}((M_1 \times M_2) \otimes_{R_1 \times R_2} R_i) \leq \text{Gpd}_{R_1 \times R_2}(M_1 \times M_2) \leq d$. This follows that $\text{G-gldim}(R_i) \leq d$ for $i = 1, 2$.

Conversely, assume that $G\text{-gldim}(R_i) \leq d$ for $i = 1, 2$ (where d is a positive integer), and consider an $R_1 \times R_2$ -module M . We may write $M = M_1 \times M_2$ where $M_i = M \otimes_{R_1 \times R_2} R_i$ for $i = 1, 2$. By hypothesis and from [2, Proposition 3.14], we may apply Lemma 3.3, and so $\text{Gpd}_{R_1 \times R_2}(M_1 \times M_2) \leq \sup\{\text{Gpd}_{R_1}(M_1), \text{Gpd}_{R_2}(M_2)\} \leq d$. Therefore, $G\text{-gldim}(R_1 \times R_2) \leq d$. ■

We can now construct a family of non-Noetherian rings $\{R_i\}_{i \geq 1}$ such that $G\text{-gldim}(R_i) = i$ and $\text{wdim}(R_i) = \infty$ for all $i \geq 1$, as follows:

Example 3.4. Consider a non-semisimple quasi-Frobenius ring R , and a non-Noetherian hereditary ring S . Then, for every positive integer n , we have: $G\text{-gldim}(R \times S[X_1, X_2, \dots, X_n]) = n + 1$ and $\text{wdim}(R \times S[X_1, X_2, \dots, X_n]) = \infty$.

Proof. Since S is a non-Noetherian hereditary ring, $\text{gldim}(S) = 1$. Then, the first equality follows immediately from Hilbert's syzygy theorem, Theorem 3.1, and since $G\text{-gldim}(R) = 0$ (since R is quasi-Frobenius). Now, if $\text{wdim}(R \times S[X_1, X_2, \dots, X_n]) < \infty$, then, from [2, Proposition 3.11], we have: $\text{gldim}(R \times S[X_1, X_2, \dots, X_n]) = G\text{-gldim}(R \times S[X_1, X_2, \dots, X_n]) = n + 1$. Thus, $\text{gldim}(R) \leq n + 1$ (by [3, Chapter VI, Exercice 10, page 123]). But, this is absurd, since R is a non-semisimple quasi-Frobenius ring. ■

Now we study the Gorenstein weak dimension case.

Theorem 3.5. Let $\{R_i\}_{i=1, \dots, m}$ be a family of coherent rings. Then:

$$G\text{-wdim}\left(\prod_{i=1}^m R_i\right) = \sup\{G\text{-wdim}(R_i), 1 \leq i \leq m\}.$$

Similarly to the Gorenstein global dimension case, we need the following lemmas:

Lemma 3.6. Let $\{R_i\}_{i=1, \dots, m}$ be a family of coherent rings such that all injective $R_i[X]$ -modules have finite flat dimension, for $i=1, \dots, m$. Let M_i be a family of R_i -module for $i=1, \dots, m$. If each M_i is a (strongly) flat R_i -module, then $\prod_{i=1}^m M_i$ is a (strongly) flat $(\prod_{i=1}^m R_i)$ -module.

Namely, we have: $\text{Gfd}_{(\prod_{i=1}^m R_i)}(\prod_{i=1}^m M_i) \leq \sup\{\text{Gfd}_{R_i}(M_i), 1 \leq i \leq m\}$.

Proof. First, note that $\prod_{i=1}^m R_i$ is also a coherent ring (by [6, Theorem 2.4.3]). Using the results of (strongly) Gorenstein flat modules ([1, Theorem 3.5], [2, Proposition 4.6], and [7, Theorem 3.7]), using the fact that, if I is an injective $R_1 \times R_2$ -module, then $I = \text{Hom}_{R_1 \times R_2}(R_1 \times R_2, I) = \text{Hom}_{R_1 \times R_2}(R_1, I) \times \text{Hom}_{R_1 \times R_2}(R_2, I)$ where each of $\text{Hom}_{R_1 \times R_2}(R_i, I)$, for $i = 1, 2$, is an injective R_i -module, and by Lemma 3.7 below, the argument is similar to the proof of Lemma 3.3. ■

Lemma 3.7. Let $R_1 \times R_2$ be a product of rings R_1 and R_2 , and let F_i be an R_i -module for $i=1, 2$. Then, $\text{fd}_{R_1 \times R_2}(F_1 \times F_2) = \sup\{\text{fd}_{R_1}(F_1), \text{fd}_{R_2}(F_2)\}$.

Proof. We have $(F_1 \times F_2) \otimes (R_1 \times 0) = (F_1 \times F_2) \otimes (R_1 \times R_2 / 0 \times R_2) = F_1 \times 0$. Then, since R_1 is a projective $R_1 \times R_2$ -module, [3, Chapter VI, Exercice 10, page 123] gives: $\text{fd}_{R_1}(F_1 \times 0) = \text{fd}_{R_1}((F_1 \times F_2) \otimes (R_1 \times 0)) \leq \text{fd}_{R_1 \times R_2}(F_1 \times F_2)$. The same: $\text{fd}_{R_2}(0 \times F_2) \leq \text{fd}_{R_1 \times R_2}(F_1 \times F_2)$.

Thus, $\sup\{\text{fd}_{R_1}(F_1), \text{fd}_{R_2}(F_2)\} \leq \text{fd}_{R_1 \times R_2}(F_1 \times F_2)$.

Conversely, from [3, Chapter VI, Exercice 10, page 123], we have:

$\text{fd}_{R_1 \times R_2}(F_1 \times 0) \leq \text{fd}_{R_1 \times 0}(F_1 \times 0) = \text{fd}_{R_1 \times 0}(F_1)$ and $\text{fd}_{R_1 \times R_2}(0 \times F_2) \leq \text{fd}_{0 \times R_2}(0 \times F_2) = \text{fd}_{0 \times R_2}(F_2)$. Therefore, $\text{fd}_{R_1 \times R_2}(F_1 \times F_2) = \sup\{\text{fd}_{R_1 \times R_2}(F_1 \times 0), \text{fd}_{R_1 \times R_2}(0 \times F_2)\} \leq \sup\{\text{fd}_{R_1}(F_1), \text{fd}_{R_2}(F_2)\}$, as desired. ■

Proof of Theorem 3.5. Using [2, Theorem 4.11], [7, Proposition 3.10], and Lemma 3.6, the argument is similar to the proof of Theorem 3.1. ■

Theorem 3.5 allows us to construct a family of non-Noetherian coherent rings $\{R_i\}_{i \geq 1}$ such that $\text{G-wdim}(R_i) = i$, $\text{G-gldim}(R_i) > i$, and $\text{wdim}(R_i) = \infty$ for all $i \geq 1$, as follows:

Example 3.8. Consider a non-semisimple quasi-Frobenius ring R and a semihereditary ring S which is not hereditary. Then, for every positive integer n , we have: $\text{G-wdim}(R \times S[X_1, X_2, \dots, X_n]) = n+1$, $\text{G-gldim}(R \times S[X_1, X_2, \dots, X_n]) > n+1$, and $\text{wdim}(R \times S[X_1, X_2, \dots, X_n]) = \infty$.

Proof. Recall, at first, that every semihereditary ring T is stably coherent, that is the polynomial ring $T[X_1, \dots, X_j]$ is coherent for every positive integer j [6, Corollary 7.3.4].

The proof of the first and the last equality is similar to the one of Example 3.4.

Now, assume, by absurd, that $\text{G-gldim}(R \times S[X_1, X_2, \dots, X_n]) \leq n+1$.

From Theorem 3.1, $\text{G-gldim}(S[X_1, X_2, \dots, X_n]) \leq n+1$. But, since S is semihereditary, $\text{wdim}(S) \leq 1$, then $\text{wdim}(S[X_1, X_2, \dots, X_n]) \leq n+1$. Thus, from [2, Proposition 3.11], we have: $\text{gldim}(S[X_1, X_2, \dots, X_n]) = \text{G-gldim}(S[X_1, X_2, \dots, X_n]) \leq n+1$. Hence, $\text{gldim}(S) \leq 1$. But, this means that S is hereditary which is absurd by hypothesis. ■

REFERENCES

- [1] D. Bennis and N. Mahdou; Strongly Gorenstein projective, injective, and flat modules, J. Pure Appl. Algebra (In Press).
- [2] D. Bennis and N. Mahdou; Global Gorenstein Dimensions, submitted for publication. Available from math.AC/0611358 v2 28 Feb 2007.
- [3] H. Cartan and S. Eilenberg; Homological Algebra, Princeton University Press, 1956.
- [4] L. W. Christensen; Gorenstein dimensions, Lecture Notes in Math., 1747, Springer, Berlin, (2000).
- [5] L. W. Christensen, A. Frankild, and H. Holm; On Gorenstein projective, injective and flat dimensions - a functorial description with applications, J. Algebra **302** (2006), 231–279.
- [6] S. Glaz; Commutative Coherent Rings, Springer-Verlag, Lecture Notes in Mathematics, 1371 (1989).
- [7] H. Holm; Gorenstein homological dimensions, J. Pure Appl. Algebra **189** (2004), 167–193.
- [8] Y. Iwanaga; On rings with finite self-injective dimension, Comm. Algebra **7** (4) (1979), 393–414.
- [9] I. Kaplansky; Commutative Rings, Allyn and Bacon, Boston, (1970).
- [10] N. Mahdou; On Costa's conjecture, Comm. Algebra **29** (7) (2001) 2775–2785.
- [11] J. J. Rotman; An introduction to homological algebra, Academic Press, New York, 1979.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES AND TECHNOLOGY, UNIVERSITY S. M. BEN ABDELLAH, FEZ 30000, MOROCCO

E-mail address: driss.bennis@hotmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES AND TECHNOLOGY, UNIVERSITY S. M. BEN ABDELLAH, FEZ 30000, MOROCCO

E-mail address: mahdou@hotmail.com